

The Richard Stockton College of New Jersey
Mathematical Mayhem 2013
Group Round

March 23, 2013

Name: _____

Name: _____

Name: _____

High School: _____

Instructions:


- This round consists of **5** problems worth **16** points each for a total of **80** points.
- Each of the 5 problems is free response.
- Write your complete solution in the space provided including all supporting work.
- No calculators are permitted.
- This round is **75 minutes** long. **Good Luck!**

OFFICIAL USE ONLY:


Problem #	1	2	3	4	5	Total
Points Earned						

• Group Round •

Problem 1. A polyomino is a contiguous shape formed by gluing together squares edge to edge. A polyomino made up of 4 squares is called a tetromino. There are 5 different tetrominoes, as shown below.




Flipping or rotating a tetromino does not make it a different tetromino. For instance, the four tetrominoes shown below are all considered to be the same tetromino.




A polyomino made up of 5 squares is called a pentomino. How many different pentominoes are there?

Solution to Question 1. There are 12 pentominoes, pictured below.



Problem 3. How many total squares are there in a 100 × 100 grid? How many total squares are there in a $n \times n$ grid? For example, there are 5 squares in the 2 × 2 grid shown below.



Solution to Question 3. The number of squares in an $n \times n$ grid is the sum

$$1^2 + 2^2 + \dots + (n-1)^2 + n^2 = \frac{n(n+1)(2n+1)}{6}.$$

For $n = 100$, this is .

Problem 4.

(A.) When I sum five numbers in every possible pair combination, I get the values:

0, 1, 2, 4, 7, 8, 9, 10, 11, 12.

What are the original 5 numbers?

(B.) When I sum a different set of five numbers in every possible group of 3, I get the values:

0, 3, 4, 8, 9, 10, 11, 12, 14, 19.

What are the original 5 numbers?

(C.) Is it possible to find a set of 5 numbers that when summed in every possible pair combination results in the sums

1, 2, 3, 4, 5, 6, 7, 8, 9, 10?

Is it possible to find a set of 5 numbers that when summed in every possible group of 3 results in those sums? For each situation, find an example or prove it's impossible.

Solution to Question 4.

(A.) The sum of the pairwise sums is 64, and this counts each of the original five numbers four times, so the sum of the original five numbers is 16. The sum of the largest two of the original five numbers is 12, and the sum of the smallest two is 0, so the middle number is 4. (i.e. $16 - 12 - 0 = 4$) The sum of the largest and middle is the second largest sum, 11, so the largest must be 7, and the second largest is 5. (i.e. $12 - 7 = 5$) In order for the second-smallest sum to be 1, one of the numbers has to be -3, so the numbers are $\boxed{3, 3, 4, 5, 7}$.

(B.) The sum of the triples is 90, and this counts each of the original five numbers six times, so the sum of the original five numbers is 15. The largest triple sum is 19, so the two not included sum to -4. The next-largest triple is 14, so the pair not included sum to 1, etc. So the pairwise sums are -4, 1, 3, 4, 5, 6, 7, 11, 12, 15. The sum of the largest two of the original five is 15, and the sum of the smallest two is -4, so the middle number is 4. The sum of the largest and middle is 12, so the largest must be 8, and the second largest must be 7. The sum of the smallest and middle is 1, so the smallest must be -3, so the numbers are $\boxed{3, -1, 4, 7, 8}$.

(C.) $\boxed{\text{No}}$, because the sum of 1-10 must be four times the sum of the five numbers, but the sum of 1-10 is 55, which is odd, and so it is not a multiple of four (as would be required for sums of pairs) or a multiple of six (as would be required for sums of triples). Moreover, any sequence of ten consecutive integers sums to an odd number, so no such sequence is possible.

